

Proceedings of the Iowa Academy of Science

Volume 46 | Annual Issue

Article 34

1939

The Construction and Use of a Concave Grating Spectrograph

W. C. Oelke
Grinnell College

George Montross
Grinnell College

Let us know how access to this document benefits you

Copyright ©1939 Iowa Academy of Science, Inc.

Follow this and additional works at: <https://scholarworks.uni.edu/pias>

Recommended Citation

Oelke, W. C. and Montross, George (1939) "The Construction and Use of a Concave Grating Spectrograph," *Proceedings of the Iowa Academy of Science*, 46(1), 197-197.

Available at: <https://scholarworks.uni.edu/pias/vol46/iss1/34>

This Research is brought to you for free and open access by the Iowa Academy of Science at UNI ScholarWorks. It has been accepted for inclusion in Proceedings of the Iowa Academy of Science by an authorized editor of UNI ScholarWorks. For more information, please contact scholarworks@uni.edu.

of moisture, but within the temperature ranges studied, we feel justified in stating that the solvating power of anhydrous magnesium bromide is not a function of its preparation temperature.

PHYSICAL CHEMISTRY LABORATORIES,
STATE UNIVERSITY OF IOWA,
IOWA CITY, IOWA.

THE CONSTRUCTION AND USE OF A CONCAVE GRATING SPECTROGRAPH

W. C. OELKE AND GEORGE MONTROSS

In order to learn some of the fundamentals of analytical spectrography, a concave grating spectrograph was constructed as a co-operative student-faculty project. Details of construction and methods of use were given. Both qualitative and quantitative applications of the instrument are considered.

DEPARTMENT OF CHEMISTRY,
GRINNELL COLLEGE,
GRINNELL, IOWA.

A QUANTITATIVE SPECTROGRAPHIC STUDY ON THE EFFECTS OF THE ALKALI METALS UPON THE DETERMINATION OF CALCIUM

LOUIS WALDBAUER AND JOHN A. MEANS

It has been known for some time that in spectrographic analysis each element present has some effect upon the other elements present. Our investigation of the effect of the alkali metals on calcium showed that, in general, the intensity of the spectral lines of calcium increased with an increase in the percentage of the alkali metal present. However, there is a decrease in the intensifying effect at the higher concentrations. The chlorides of the alkali metals used were in concentrations varying from 1.0 per cent to 0.031 per cent, and the concentrations of the calcium solutions used varied approximately over the same range. Slides were